

Algebraic Models: Coordinates, Scales, and Dynamical Symmetries

Michael W.N. Ibrahim

*Center for Theoretical Physics, Sloane Laboratory,
Yale University, New Haven, Connecticut 06520-8120*

Abstract

We discuss the variety of coordinates often used to characterize the coherent state classical limit of an algebraic model. We show selection of appropriate coordinates naturally motivates a procedure to generate a single particle Schrödinger hamiltonian which, for low energy states, gives equivalent results to a bosonic algebraic model to leading order in N . The process is used to study the associated geometries of the dynamical symmetries of $U(3)$. By demanding that the inner product be preserved in the Schrödinger picture we conclude that different dynamical symmetries correspond to different scales.

1 Introduction


Recently, hybrid algebraic-Schrödinger approaches for the study of transition intensities in molecules have been introduced [1–3]. In these approaches one relies on Schrödinger/algebraic correspondences to introduce operators in the algebra which are functions of configuration space parameters. For the $U(2)$ model considered in [1,2] one does so by relying on correspondences between a particular dynamical symmetry and a Schrödinger Hamiltonian [4]. For more complicated algebras such correspondences are not available. Further, the inner-product of two algebraic states is not equal to the Schrödinger inner-product of their images. Since, in some circumstances, the inner-product of the algebra has a natural interpretation as a Schrödinger overlap [5] such shortcomings are an impediment to understanding molecular structure.

It would therefore be useful to have a generic correspondence—independent of a particular dynamical symmetry—for the interpretation of algebraic parameters as geometric quantities which reside in configuration space (which is noticeably absent in the algebraic models). We provide such an interpretation by exploiting the many approximate correspondences between algebraic and Schrödinger pictures. We demonstrate how one may turn an algebraic hamiltonian into a traditional Schrödinger (traditional meaning kinetic plus potential—with no coordinate dependent mass terms) single particle hamiltonian which will give the same results to leading order in N , the label for the symmetric representation of the algebra. Our procedure is most similar to the re-quantization used by [6] to study

spectra. We proceed further to study the wavefunctions and inner product. Our approach differs from other studies which have examined the correspondence between algebraic and Schrödinger inner products [7] since they rely on particular dynamical symmetries. We work out our procedure explicitly for $U(3)$ and conclude that different chains correspond not only to different geometries but different scales.

2 Algebraic Models: Constructing a Connection with Configuration Space

Numerous correspondences between algebraic and Schrödinger methods and their classical limits can be succinctly summarized in the following simple diagram:



Hamiltonians on the right hand side are written in terms of geometric parameters whereas those on the left are written in terms of algebraic ones. Thus, a careful exploitation of the horizontal correspondences should relate the different parameters. We propose to do so by travelling counter-clockwise around the diagram—re-quantizing the algebraic model.

The vertical correspondences are extremely well defined and have been thoroughly discussed in the literature [6,8–11]. The relationship between the $U(n) \supset U(n-1)$ and the simple harmonic oscillator (SHO) has also been discussed [12] and can be formalized by considering the contraction limit of the algebra [13,3]. Similarly, if one finds appropriate coordinates in a patch of the coherent state phase space (CSPA) one may embed this region within a standard phase space and have a perfect copy of the dynamics for local trajectories [14].

The horizontal correspondences are approximate on the whole given that the phase spaces are topologically different (compact versus non-compact) in the classical regimes and one has a finite versus infinite dimensional Hilbert space in the corresponding quantum cases. However, with careful selection of coordinates it will be seen that these incompatibilities will not effect the lower bound states in a re-quantization scheme.

It is to the selection of these coordinates we now turn. We must find coordinates in a patch of CSPA which behave like positions and momenta and then identify these as coordinates covering the entire canonical phase space. In the language of geometry, we wish

to find coordinates in which the symplectic form determining the dynamics is in Darboux form: $\omega = dp \wedge dq$ [14]. Historically, it has been useful to think of the imaginary part of the coherent state parameter α as a momenta [12,6]. We adopt that convention here to eliminate any remaining coordinate choice ambiguity.

Examining the action in the propagator path integral [11,6]: one finds that the classical hamiltonian is given by the coherent state expectation, $H = \langle \alpha | \hat{H} | \alpha \rangle$; $|\alpha\rangle = \frac{1}{\sqrt{N!}}(\sqrt{1 - \alpha \cdot \alpha^*} \sigma^\dagger + \alpha \cdot \tau^\dagger)^N |0\rangle$ denotes the coherent state in group coordinates [8]; σ and τ are typically scalar and tensor operators of $O(3)$ respectively; and the hamiltonian evolution is determined by the symplectic form $\omega = i\hbar d\langle \alpha | d | \alpha \rangle$ which in group coordinates is explicitly:

$$\omega = Ni\hbar d\alpha_\mu^* \wedge d\alpha_\mu. \quad (2.2)$$

Deviating slightly from the typical development we let

$$\alpha_\mu = \sqrt{\frac{\overline{m\omega}}{2}} \left(\frac{q_\mu}{\sqrt{N}} + \frac{i}{\overline{m\omega}} \frac{p_\mu}{\sqrt{N}} \right), \quad (2.3)$$

where q and p are now dimensionful quantities and $\overline{m\omega}$ has units of length over momentum and has the physical interpretation of the ratio of the natural distance and momenta scales of the problem (see appendix A). In these coordinates the symplectic two-form is in standard Darboux form $\omega = dp_\mu \wedge dq_\mu$, or equivalently $\{q, p\}_{\text{PB}} = 1$.

We have been extremely explicit in order to contrast our choice with those of the literature. For instance in the coordinates of [6] $\{\tilde{q}, \tilde{p}\}_{\text{PB}} = N$. This point is obfuscated because the correct equations of motion are maintained by dividing the classical action (and hence ω) by N . Such a procedure, however, leads to improper quantization. Similarly in terms of the projective coordinates [8] used to establish algebraic-geometric correspondences in such references as [15,12] one has $\{\tilde{q}, \tilde{p}\}_{\text{PB}} = N(1 + \tilde{q}^2 + \tilde{p}^2)^2$. Thus only near the domain $\tilde{p}^2 + \tilde{q}^2 \approx 0$ is it suitable to interpret these coordinates as the natural coordinates of a cotangent bundle over configuration space.

Classical Hamiltonians obtained from our coordinates usually have contributions such as p_r^4 and $p_r^2 r^2$. This leads to canonical quantization ordering ambiguities as well as higher degree differential equations. However, if we focus on the lowest bound states we may use an approximate hamiltonian which is valid in the classical regions whose paths most greatly contribute to these states. In the spirit of the stationary phase approximation, the appropriate region would be near the fixed point of the hamiltonian flow, i.e. where $dH = 0$.

Fortunately, since the CSPS limit of algebraic hamiltonians typically have a convenient momenta dependence, i.e. $H = p_r^2 f(p_r^2, r^2) + V_{\text{eff}}(r)$, this condition implies we are looking in the region of reduced phase space near $p_r = 0$ and $r = r^*$ where $V'_{\text{eff}}(r^*) = 0$. These conditions are eminently reasonable.

Near this region the dynamics can be given by approximating the hamiltonian:

$$H(p_r, r) \approx H|_{p_r=0, r=r^*} + \frac{1}{2} \left(V''_{\text{eff}}(r^*)(r - r^*)^2 + H_{1^2}|_{p=0, r=r^*} p_r^2 \right), \quad (2.4)$$

where all other lower order terms vanish. Hence, at least locally around the fixed point of the system, our ordering ambiguities are resolved. If r^* is sufficiently far from the $r = 0$ phase space boundary one can sensically re-quantize.

Given our coordinate choice each higher derivative in the Taylor expansion of H in equation 2.4 will be down by a factor of $\frac{1}{\sqrt{N}}$. Thus this approximation may be considered as an expansion in large N . Note, however that this procedure is distinct from other ‘large N ’ techniques such as the contraction limit [13, 3]. Indeed, we will see in section 3.2 that this procedure, in the specific instance for Hamiltonians with the dynamical symmetry $U(n) \supset U(n-1)$, essentially reproduces the contraction limit results.

We have carried out this discussion in a reduced phase space with a radial coordinate. That is, due to $O(3)$ invariance the fixed point is not a true minima in the global phase space. In instances where one has a true fixed point in the global phase space (e.g. when the minima is at $r = 0$) a similar procedure applies.

The introduction of the scales in equation 2.3 appears arbitray. However, since we desire a generic correspondence which preserves the inner product structure of the algebra in the Schrödinger picture, once a scale is picked for one Hamiltonian it must be fixed for others, i.e. any scale change would influence the overlap of eigenstates which should be equated with the inner product of the representation. Thus, the scale for each hamiltonian must be functionally related to the scale of another. If the relationship is not trivial $\overline{m\omega}$ may depend on the parameters within the hamiltonian. This dependence of the scale ratio, $\overline{m\omega}$, will be suppressed until such subtleties must be considered, at which time we will supplement our notation with subscripts.

3 An Explicit Example: $U(3)$

3.1 Coherent State Limit

The algebraic approach for 2D problems was presented by [16]. One considers symmetric representations of $u(3)$ realized with the two chains of interest:

$$\begin{aligned} U(3) \supset U(2) \supset O(2) & \quad \text{I} \\ U(3) \supset O(3) \supset O(2) & \quad \text{II} \end{aligned} \quad (3.1)$$

We use the same notation as [16] for the generators. However, we select a different $O(3)$ subgroup which is generated by $\hat{R}_+$, $\hat{R}_-$, and $\hat{l}$. Our choice introduces more ‘coordinate like’ terms of the classical Hamiltonian. The choice makes no difference for spectra and only an overall phase for FC overlaps [17]. Please note that the $O(3)$ group is a dynamical symmetry subgroup and does not have the interpretation of a rotation in configuration space.

The general Hamiltonian of a $U(3)$ model is

$$H = \epsilon \hat{n} + \delta \hat{n}(\hat{n} + 1) + \beta \hat{l}^2 - A \hat{W}^2, \quad (3.2)$$

where ϵ , δ , and A are taken as positive or 0. Setting $A = 0$ ($\epsilon = \delta = 0$) gives a Hamiltonian with dynamical symmetry I (II). The basis corresponding to chain I is labeled by the

eigenvalues of $\hat{n}$ and $\hat{l}$ respectively. The basis corresponding to chain II is labeled by the eigenvalues of $\hat{W}^2 = \frac{1}{2}(\hat{R}_+\hat{R}_- + \hat{R}_-\hat{R}_+) + \hat{l}^2$ and again $\hat{l}$.

Taking the appropriate expectations one may calculate the classical limits of the various operators. The results are enumerated in Table 1.

We reduce the phase space in the natural way:

$$\begin{aligned} q_x &= r \cos \theta & q_y &= r \sin \theta \\ p_x &= p_r \cos \theta - p_\theta r \sin \theta & p_y &= p_r \sin \theta + p_\theta r \cos \theta, \end{aligned} \quad (3.3)$$

implying

$$q \cdot q = r^2, \quad p \cdot p = p_r^2 + \frac{l_{\text{cl}}^2}{r^2}, \quad (3.4)$$

where $l_{\text{cl}}^2 = p_\theta^2 r^4$ is a constant of the motion.

3.2 $U(2)$ Chain

We work out the re-quantization procedure in the case where $A = \beta = \delta = 0$. In this instance we have the Hamiltonian on the reduced phase space:

$$H_{\text{cl}} = \epsilon \frac{\overline{m\omega}}{2\hbar} \left(\frac{1}{(\overline{m\omega})^2} (p_r^2 + \frac{l_{\text{cl}}^2}{r^2}) + r^2 \right). \quad (3.5)$$

The canonical quantization scheme yields the following prescription:

$$p_r^2 \mapsto -\frac{\hbar^2}{r} \frac{d}{dr} \left(r \frac{d}{dr} (\cdot) \right), \quad l \mapsto \hbar l, \quad (3.6)$$

where l now labels irreps. of $SO(2)$. This yields the quantum Hamiltonian:

$$H = \epsilon \frac{\hbar}{2\overline{m\omega}} \left(-\frac{1}{r} \frac{d}{dr} \left(r \frac{d}{dr} (\cdot) \right) + \frac{l^2}{r^2} + \frac{(\overline{m\omega})^2}{\hbar^2} r^2 \right), \quad (3.7)$$

which we immediately recognize as a multiple of a 2D circular oscillator with energy levels spaced by $\Delta E = \epsilon$ [18]. This result perfectly agrees with direct algebraic evaluation.

We proceed similarly for δ different from 0. The classical Hamiltonian is then of the form:

$$H = N(\epsilon + 2\delta)n_{\text{cl}} + \delta N(N-1)(n_{\text{cl}})^2. \quad (3.8)$$

In Cartesian coordinates the conditions for a fixed point are:

$$\begin{aligned} dH = 0 &= N[\epsilon + 2\delta + 2\delta(N-1)n_{\text{cl}}] \\ &\times \left(\frac{\partial n_{\text{cl}}}{\partial x} dx + \frac{\partial n_{\text{cl}}}{\partial y} dy + \frac{\partial n_{\text{cl}}}{\partial p_x} dp_x + \frac{\partial n_{\text{cl}}}{\partial p_y} dp_y \right), \end{aligned} \quad (3.9)$$

which is true at $x = y = p_x = p_y = 0$ (hence our use of Cartesian as opposed to radial coordinates). Near this point to second order our Hamiltonian behaves like:

$$H \approx N(\epsilon + 2\delta)n_{\text{cl}}. \quad (3.10)$$

Re-quantization proceeds exactly as in the previous case leading to an energy spacing of $\Delta E \approx \epsilon + 2\delta$. The corresponding algebraic Hamiltonian ($H = \epsilon\hat{n} + \delta\hat{n}(\hat{n} + 1)$) has spacing for the lowest levels of $\Delta E = \epsilon + 2\delta$.

Additionally the same low energy eigenfunctions diagonalize both $U(2)$ Hamiltonians in both the algebraic and approximate re-quantizing prescriptions (i.e. in both prescriptions the inner-product of the lowest eigenfunctions of the Hamiltonian with $\delta = 0$ and $\delta \neq 0$ is 1).

3.3 Radially Displaced Oscillators—Near the $O(3)$ Chain

In the situation where our Hamiltonian is near the $O(3)$ dynamical symmetry the fixed point of the Hamiltonian flow is not at $r = 0$.

The potential is given by:

$$V_{\text{eff}}(r) = \frac{\overline{m\omega}}{\hbar} \left[(X - Z) \frac{1}{2} r^2 + (Y + Z) \frac{\overline{m\omega}}{4N\hbar} r^4 + X \frac{1}{2} \frac{1}{(\overline{m\omega})^2} \frac{l^2}{r^2} + Y \frac{1}{4N\hbar(\overline{m\omega})^3} \frac{l^4}{r^4} \right], \quad (3.11)$$

where we have made the definitions $X = \epsilon + 2\delta + \beta$, $Y = \delta(N - 1)$, and $Z = 4A(N - 1)$. The physically interesting regime [5] is ‘near’ the $O(3)$ chain, i.e. $\frac{X}{Z}$ and $\frac{Y}{Z}$ are both small. In this limit it is easy to find a perturbative solution for the condition $V'_{\text{eff}}(r^*) = 0$:

$$(r^*)^2 \approx \frac{N\hbar}{\overline{m\omega}} \left(1 + \frac{X}{Z} \left(\frac{l^2}{(N\hbar)^2} - 1 \right) - \frac{Y}{Z} \right), \quad (3.12)$$

where now it is clear that the presumption that r^* is large depends on N being large. Note that having X or Y different from 0 decreases r^* . That is, the $O(3)$ chain represents the maximum possible displacement. r^* may increase with increasing l however, corresponding to a larger angular momentum barrier.

Since l can be as large as N we regard l^2/N^2 as an independent quantity—even though in practice it is of the same order as terms which go $1/N^2$. In anticipation of this we dropped the term going with $\frac{l^2}{N^4}$ for simplicity.

When one moves slightly off the $O(3)$ chain different l subspaces which were degenerate split resulting in the lowest energy difference, ΔE , having a slight l dependence. As we’ve seen this dependence has entered in lowest order in the form of terms $\frac{l^2}{N^2}$. Since the ratio $\frac{l}{N}$ can be regarded as a separate independent small quantity ignoring terms like $O(\frac{1}{\sqrt{N}})$ does not mean we lose all l dependence in our calculation. Our results for the energy level spacing will only be correct to leading order in N . The l space splitting behavior will be given correctly—but the actual values of the splitting will be insignificant compared to other (non- l dependent) contributions we have ignored for smaller l .

Continuing we see:

$$V''_{\text{eff}}(r^*) \approx 2Z \frac{\overline{m\omega}}{\hbar} \left[1 + \frac{X}{Z} \left(3 \frac{l^2}{(N\hbar)^2} - 1 \right) \right], \quad (3.13)$$

$$H_{p_r^2}|_{p_r=0, r=r^*} \approx \frac{Z}{2\hbar(\overline{m\omega})} \left[1 + \frac{X}{Z} \left(\frac{l^2}{(N\hbar)^2} + 1 \right) + \frac{Y}{Z} \left(2 \frac{l^2}{(N\hbar)^2} + 1 \right) \right]. \quad (3.14)$$

Near the fixed point the Hamiltonian's behavior is given by equation 2.4. However, the radial quantum mechanics depends crucially on the point $r = 0$. Since equation 2.4 does not reproduce the original Hamiltonian's behavior near this point the approximation becomes suspect. It would be more reasonable to quantize:

$$\frac{1}{2}H_{p_r^2}|_{p=0, r=r^*}p_r^2 + \tilde{V}(r), \quad (3.15)$$

where $\tilde{V}(r)$ has the same behavior near the fixed point as the approximate Hamiltonian and the behavior near $r = 0$ of the original Hamiltonian. That is:

$$\lim_{r \rightarrow 0} r^4 \tilde{V}(r) = \frac{Y}{4} \frac{l^4}{N^2 \hbar^2 (\overline{m\omega})^2} \quad (3.16)$$

$$\tilde{V}(r) = H|_{p_r=0, r=r^*} + \frac{1}{2}V''_{\text{eff}}(r^*)(r - r^*)^2 + O((r - r^*)^3). \quad (3.17)$$

Additionally, of course, $\tilde{V}$ should not introduce any other minima at other locations.

Following the prescription for the harmonic oscillator we use the radial quantization scheme (equation 3.6), substituting for the wavefunction, $\Psi = \frac{\psi}{\sqrt{r}}$, we obtain:

$$-H_{p_r^2} \frac{\hbar^2}{2} \psi'' + \left[\tilde{V}(r) - \frac{H_{p_r^2} \hbar^2}{8r^2} \right] \psi = E\psi, \quad (3.18)$$

where $\psi(0) = 0$ and $\psi(r \rightarrow \infty) = 0$. The form of this equation is that of a 1-D Schrödinger equation except the left boundary condition is applied at $r = 0$. At this point the $r \approx 0$ dependence of $\tilde{V}$ becomes crucial. Since $\tilde{V}$ blows up more quickly than the counter term blows down the sum of the two still has a minima, although it should be shifted inward from r^* . Calling the new minima $r^{**} = r^*(1 - \epsilon)$ and using our knowledge of the behavior of $\tilde{V}$ near r^* we find that the condition $\frac{d}{dr}|_{r=r^{**}} \left[\tilde{V}(r) - \frac{H_{p_r^2} \hbar^2}{8r^2} \right] = 0$ yields:

$$\epsilon \approx \frac{1}{16N^2 - 3} \left[1 - \frac{16N^2}{16N^2 - 3} \left(4 \frac{X}{Z} \left(\frac{l^2}{N^2} - 1 \right) - \frac{Y}{Z} \left(2 \frac{l^2}{N^2} + 3 \right) \right) \right] \quad (3.19)$$

This can be expanded in N with leading contribution $O(\frac{1}{N^2})$. Recalling that N essentially counts the number of bound states (hence effectively measures the depth of the potential) it is reasonable that $\frac{1}{N^2}$ should parameterize ϵ . Of course, since we are only working in leading order in N we have $\epsilon \approx 0$ and consequently $r^{**} \approx r^*$. Our primary intent of displaying equation 3.19 is that the condition of N being large is now clarified. Since the prefactor must be small we find $2N \gg 1$.

Returning to equation 3.18 we notice that in our approximation (r^* or r^{**} is large) a solution with the left boundary condition at $r = -\infty$ will, to a good approximation, satisfy the boundary condition at $r = 0$. In this instance the lower eigenvalues are given by $E \approx \hbar\omega$ where

$$\omega^2 = H_{p_r^2} \left[\tilde{V}''(r^{**}) - \frac{3H_{p_r^2} \hbar^2}{4(r^{**})^4} \right]. \quad (3.20)$$

Given the properties of $\tilde{V}$ we have $\tilde{V}''(r^{**}) \approx \tilde{V}''(r^*) = V_{\text{eff}}''(r^*)$. The second term is $O(\frac{1}{N^2})$ and may be ignored.

Thus we have

$$\Delta E \approx Z \left[1 + 2 \frac{X}{Z} \frac{l^2}{N^2} + \frac{1}{2} \frac{Y}{Z} \left(2 \frac{l^2}{N^2} + 1 \right) \right]. \quad (3.21)$$

We see that on the $O(3)$ chain ($X = Y = 0$) we have a spacing of $\Delta E \approx Z \approx 4AN$, as compared to the exact algebraic expression:

$$\Delta E = A [N(N+1) - (N-2)(N-1)] = A(4N-2), \quad (3.22)$$

which agrees to leading order in N as advertised.

Next we compute the induced harmonic dilatation constant of the geometry (see Appendix A):

$$\alpha^2 \approx \frac{\omega}{\hbar H_{1^2}} \approx 2 \frac{(\overline{m\omega})}{\hbar} \left[1 + \frac{X}{Z} \left(\frac{l^2}{N^2} - 1 \right) - \frac{Y}{Z} \left(\frac{l^2}{N^2} + \frac{1}{2} \right) \right]. \quad (3.23)$$

Note, that the ‘concavity’ corrections depend explicitly on the algebraic parameters, indicative of the fact that our calculations have established relationships between algebraic Hamiltonians and the geometry of configuration space.

Equations 3.23 and 3.12 for the harmonic dilatation and radial displacement can be easily related to experimental data and provide useful assistance when analyzing transition intensities as to be detailed in the upcoming publication [5].

3.4 Near $O(3)$ to $O(3)$ Inner Product

We wish to calculate the inner-product of eigenstates of a hamiltonian corresponding to slightly off $O(3)$ symmetry to one on $O(3)$. In either case the approximate wavefunctions are:

$$\psi_0^{\text{disp}} \approx \frac{\sqrt{\alpha}}{\pi^{\frac{1}{4}}} \frac{e^{-\frac{1}{2}[\alpha(r-r^*)]^2}}{\sqrt{r}}, \quad (3.24)$$

where we have normalized it on the interval $[-\infty, \infty]$ (as opposed to $[0, \infty]$) as this only introduces errors of order $\frac{1}{N}$. The parameters α and r^* are determined by 3.23 and 3.12. We use α' and $r^{*'}$ as the parameters of the wavefunction exactly on the $O(3)$ chain (determined by the same equations except $X = Y = 0$). Note that there is an l dependence hidden in both r^* and α .

Since both wavefunctions are essentially 0 for $r < 0$ we may again calculate the overlap on the interval $[-\infty, \infty]$. Thus we wish to evaluate the integral

$$I_{0,0} \approx \int_{-\infty}^{\infty} \phi(\alpha; r - r^*) \phi(\alpha'; r - r^{*'}) dr, \quad (3.25)$$

where ϕ is the ground state wavefunction of a 1D SHO. The integral is easily evaluated to [1]:

$$I_{0,0} \approx \left[\frac{2\alpha\alpha'}{\alpha^2 + \alpha'^2} \right]^{\frac{1}{2}} \times \exp \left[-\frac{(\alpha\alpha'(r^{*'} - r^*))^2}{2(\alpha^2 + \alpha'^2)} \right]. \quad (3.26)$$

As stated previously the ratio of scales $\overline{m\omega}$ need not be the same in both scenarios and may indeed depend on the value of X, Y , and Z . However, this dependance must be smooth. Since the difference between these two scenarios is perturbative one concludes that $\frac{\overline{m\omega}}{m\omega'} = 1 + \delta$ where $\delta \sim O(X/Z, Y/Z)$. Given this, we have:

$$I_{0,0} \approx 1 + O\left(\left(\frac{X}{Z}\right)^2, \left(\frac{Y}{Z}\right)^2\right) \quad (3.27)$$

This is the exact overlap one would expect from perturbation theory of the algebraic model as spelled out in Appendix B.

3.5 $O(3)$ to $U(2)$ Inner-Products

Given equation 3.27, to our order the overlaps from a hamiltonian near $O(3)$ to the $U(2)$ chain should be exactly the same as the exact $O(3)$ - $U(2)$ factors.

For the $U(2)$ chain we have the ground state wavefunction [18]:

$$\psi_0^{U(2)} = \sqrt{\frac{2}{|l|!}} (\alpha_{U(2)})^{|l|+1} r^{|l|} e^{-\frac{1}{2}[\alpha_{U(2)} r]^2}, \quad (3.28)$$

where $(\alpha_{U(2)})^2 = \frac{\overline{m\omega}_{U(2)}}{\hbar}$.

Thus our overlap becomes:

$$I_{0,0}\left(\frac{\alpha_{U(2)}}{\alpha}, \alpha r^*\right) = \sqrt{\frac{2\alpha}{|l|!}} \frac{(\alpha_{U(2)})^{|l|+1}}{\pi^{\frac{1}{4}}} \exp\left[-\frac{1}{2} \frac{\alpha^2 \alpha_{U(2)}^2}{\alpha_+^2} r^{*2}\right] \times \int_0^\infty dr r^{|l|+\frac{1}{2}} \exp\left[-\frac{1}{2}(\alpha_+ r - \frac{\alpha^2 r^*}{\alpha_+})^2\right] \quad (3.29)$$

where $\alpha_+^2 = \alpha_{U(2)}^2 + \alpha^2$, and α is again defined by 3.23 with $X = Y = 0$. The integral may be expressed in terms of confluent hypergeometric functions:

$$\begin{aligned} \int_0^\infty dr r^{|l|+\frac{1}{2}} \exp\left[-\frac{1}{2}(\alpha_+ r - \frac{\alpha^2 r^*}{\alpha_+})^2\right] &= 2^{\frac{2l-1}{4}} \frac{1}{\alpha_+^{\frac{3+2l}{2}}} \exp\left[-\frac{1}{2} \left(\frac{\alpha^2 r^*}{\alpha_+}\right)^2\right] \\ &\times \left\{ \Gamma\left(\frac{3+2l}{4}\right) {}_1F_1\left(\frac{3+2l}{4}, \frac{1}{2}; \frac{1}{2} \left(\frac{\alpha^2 r^*}{\alpha_+}\right)^2\right) \right. \\ &\quad \left. + \sqrt{2} \left(\frac{\alpha^2 r^*}{\alpha_+}\right) \Gamma\left(\frac{5+2l}{4}\right) {}_1F_1\left(\frac{5+2l}{4}, \frac{1}{2}; \frac{1}{2} \left(\frac{\alpha^2 r^*}{\alpha_+}\right)^2\right) \right\}. \end{aligned} \quad (3.30)$$

We note that $\frac{\alpha^2 r^*}{\alpha_+} \sim N$ and hence use an asymptotic expansion for the hypergeometric functions [19]. Keeping only the most dominant contribution of the integral we find:

$$I_{0,0} \approx \frac{\pi^{\frac{1}{4}}}{\sqrt{|l|!}} \left(\frac{\alpha_{U(2)}}{\alpha}\right)^{|l|+1} \left(\frac{\alpha}{\alpha_+}\right)^{4+2|l|} \exp\left[-\frac{1}{2} \frac{\alpha^2 \alpha_{U(2)}^2}{\alpha_+^2} r^{*2}\right] (\alpha r^*)^{\frac{5+2|l|}{2}}. \quad (3.31)$$

Substituting the expressions for the harmonic scales, and letting $\zeta = \frac{\overline{m\omega}_{U(2)}}{\overline{m\omega}_{O(3)}}$ we find

$$I_{0,0} \approx \frac{\pi^{\frac{1}{4}}}{\sqrt{|l|!}} \left(\frac{\zeta}{2}\right)^{|l|+1} \left(\frac{2}{\zeta+2}\right)^{2+|l|} \exp\left[-N \frac{1}{1+\frac{2}{\zeta}}\right] (2N)^{\frac{5+2|l|}{4}}. \quad (3.32)$$

3.6 Scale Changes

Comparing the analytic expression of the $U(2)/O(3)$ overlaps (equation 3.32 calculated by approximately requantizing) and the expression from appendix equation B.5 (calculated by direct algebraic means) fixes the scale dependance. Comparing the exponential dependance of the two expressions,

$$e^{-\frac{\ln 2}{2}N} \text{ vs. } e^{\left[-N \frac{1}{1+\frac{2}{\zeta}}\right]}, \quad (3.33)$$

leads to the conclusion that:

$$\zeta = \frac{\overline{m\omega}_{U(2)}}{\overline{m\omega}_{O(3)}} \approx \frac{2 \ln 2}{2 - \ln 2}. \quad (3.34)$$

The rest of the dependance can be fixed by allowing ζ to have $\frac{\log N}{N}$ and $\frac{1}{N}$ corrections. To see this we let:

$$\zeta \approx \frac{2 \log 2}{2 - \log 2} \left(1 - \frac{4}{\log 2(2 - \log 2)} \frac{\log \gamma}{N}\right), \quad (3.35)$$

or equivalently

$$\left(1 + \frac{2}{\zeta}\right)^{-1} \approx \frac{\log 2}{2} - \frac{\log \gamma}{N}. \quad (3.36)$$

Substituting into the expression 3.32 we find the first order term of the large N asymptotic:

$$I_{0,0} \approx \frac{\pi^{\frac{1}{4}}}{\sqrt{|l|!}} \left[\frac{\log 2}{2}\right]^{|l|+1} \left[\frac{2 - \log 2}{2}\right] (2N)^{\frac{5+2|l|}{4}} e^{-N \frac{\log 2}{2}} \gamma. \quad (3.37)$$

Comparing with the algebraic expression we conclude that

$$\gamma = \frac{2^{\frac{3}{4}}}{2 - \log 2} (\log 2)^{-(|l|+1)} N^{-\frac{3}{2}} \quad (3.38)$$

That is, in order for the Schrödinger and algebraic prescriptions to be commensurate for the leading asymptotic in N , we must have the first orders of the asymptotics of ζ :

$$\zeta \approx \frac{2 \log 2}{2 - \log 2} \left[1 + \frac{4}{\log 2(2 - \log 2)} \left\{\frac{3 \log N}{2N} + (|l| \log \log 2 + c) \frac{1}{N}\right\}\right], \quad (3.39)$$

$$c = \log \log 2 - \log \frac{2^{\frac{3}{4}}}{2 - \log 2} \quad (3.40)$$

Calculating the inner product of the eigenstates of an algebraic hamiltonian on the $U(2)$ chain with the eigenstates of another hamiltonian off of the $U(2)$ chain is not simply analogous to the overlap of radially displaced oscillators, but analagous to the matrix elements of an operator which radially displaces *and* dilatates (much like the operator matrix elements calculated in [1]) changing the natural scale of the problem. The degree of dilatation depends on the proximity of the second hamiltonian to either chain. For chains near $U(2)$ the dilatation paramater is essentially 1. As one moves nearer to the $O(3)$ chain the dilatation parameter increases, approaching the value given by equation 3.39. This has considerable consequences for hybrid algebraic-Schrödinger analysis of molecules [5].

4 Conclusions

We have provided a procedure, via requantization, to convert a hamiltonian of an algebraic model into a Schrödinger hamiltonian which will give the same results for the lower states to leading order in N . The procedure is optimized for algebras which are interpreted as single particle hamiltonians. Thus, it seems most applicable to molecular models such as the vibron model $U(4)$ [21], the anharmonic oscillator $U(2)$ [4], and the two dimensional $U(3)$ model considered here. Although the prescription should generate Schrödinger hamiltonians in other models it remains to be seen whether or not such results would have appropriate many-body interpretations.

We have carried out the requantization process in detail for the limiting cases of the $U(3)$ model. By demanding that the wavefunction overlap of the requantized system agree with the inner product of the algebraic model we have demonstrated that different chains of the $U(3)$ model correspond to not only different geometries but different scales.

5 Acknowledgements

This work was performed in part under the U.S. Department of Energy, Contract No. DE-FG02-91ER40608. I extend my deepest gratitude to my advisor, Prof. Franco Iachello, for introducing me to the problem and his suggestions and feedback on this work. I thank Prof. Dimitri Kusnezov for also making suggestions on this manuscript. Finally, I would like to thank the Department of Energy's Institute for Nuclear Theory at the University of Washington for its hospitality during the completion of this work.

Appendices

A Scales in the Schrödinger Picture

Consider a traditional Schrödinger Hamiltonian (1-D for simplicity),

$$H = -\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + V_0 f(\bar{\alpha}x), \quad (\text{A.1})$$

with the distance scale $\bar{\alpha}$ and momenta scale given by $\frac{\bar{\alpha}}{m\omega} = (mV_0)^{-\frac{1}{2}}$.

We on occasion refer to the induced harmonic dilatation constant—which we define as the distance scale (dilatation constant of the wavefunctions) one obtains by approximating the potential about a minimum to second order. For a SHO $f(x) = \frac{1}{2}x^2$ and $V_0 = \frac{m\omega^2}{\bar{\alpha}^2}$. In this instance $\overline{m\omega} = m\omega$ and we define the distance scale $\bar{\alpha}^2 = \frac{m\omega}{\hbar}$. If x^* denotes the minimum of a more complicated potential we see that the distance scale associated with its harmonic approximation is

$$\alpha^2 = \frac{\overline{m\omega}}{\hbar} \sqrt{f''(\bar{\alpha}x^*)}. \quad (\text{A.2})$$

The induced scale depends on two quantities: (1) the ratio of the true distance and momenta scales; (2) the concavity of the potential about the minima.

B Algebraic Calculations of Overlaps near $O(3)$

We wish to find the overlaps of eigenstates of the hamiltonian

$$H = A \left(\frac{(\epsilon + \delta)}{A} \hat{n} + \frac{\delta}{A} \hat{n}(\hat{n} + 1) + \frac{\beta}{A} \hat{l}^2 - \hat{W}^2 \right) \quad (\text{B.1})$$

to the hamiltonian with $\epsilon = \delta = \beta = 0$ in the limit where A is large. We may work within subspaces of constant l so that there are no degeneracies.

We denote the basis from the $O(3)$ chain by $|[N], \omega, l\rangle$ where $\omega(\omega + 1)$ is the eigenvalue of $\hat{W}^2$. The $U(2)$ chain basis is denoted by $|[N], n, l\rangle$ analogously. We denote the transformation matrix between them by ζ , i.e. $|[N], \omega\rangle = \sum_n \zeta_n^\omega |[N], n\rangle$, where we have suppressed the implicit l dependence.

The wavefunction for the ground state may be perturbatively calculated to first order:

$$|[N], E_{\text{low}}\rangle \approx |[N], \omega = N\rangle + \sum_{\omega' \neq N} |[N], \omega'\rangle \frac{\sum_n (\zeta_n^{\omega'})^* \zeta_n^{\omega=N} [(\epsilon + 2\delta)n + \delta n^2]}{AN(N+1) - A\omega'(\omega'+1)}. \quad (\text{B.2})$$

As always in perturbation theory, to first order in $\frac{1}{A}$ the wavefunctions only have corrections orthogonal to the unperturbed wavefunction. Consequently the overlap from a slight deformation off the $O(3)$ chain to the $O(3)$ chain is identically 1 to this order.

We now concentrate on the overlap between the $O(3)$ and $U(2)$ chains. In this limit we have:

$$I_{0,0} = \langle [N], n = |l|, l | [N], \omega = N, l \rangle = \zeta_{n=|l|}^{\omega=N}. \quad (\text{B.3})$$

These coefficients are calculated in [17]:

$$\zeta_{n=|l|}^{\omega=N} = \left[\frac{(N + |l|)!}{2^{|l|} |l|! (2N - 1)!!} \right]^{\frac{1}{2}}. \quad (\text{B.4})$$

Using $(2N - 1)!! = 2^N \Gamma(N + \frac{1}{2}) / \Gamma(\frac{1}{2})$ and the Stirling approximation $\Gamma(z + b) \sim \sqrt{2\pi} e^{-z} z^{z+b-\frac{1}{2}}$ [19] this formula may be approximated for large N by:

$$I_{0,0} \approx \frac{\pi^{\frac{1}{4}}}{2^{\frac{|l|}{2}} \sqrt{|l|!}} N^{\frac{|l|}{2} - \frac{1}{4}} e^{-\frac{\log 2}{2} N}. \quad (\text{B.5})$$

If one wishes to algebraically calculate FC transitions to higher energy $U(2)$ bound states one may proceed in the exact same fashion to find

$$I_{n,0} \approx \frac{\pi^{\frac{1}{4}} N^{\frac{n}{2} - \frac{1}{4}} e^{-\frac{\log 2}{2} N}}{2^{\frac{n}{2}} \sqrt{(\frac{n+l}{2})! (\frac{n-l}{2})!}}. \quad (\text{B.6})$$

References

- [1] F. Iachello, M. Ibrahim, J. Phys. Chem. A., **102**, 9427 (1998).
- [2] T. Müller, P. Dupré, P.H. Vaccaro, F. Pérez-Bernal, M. Ibrahim, F. Iachello, Chemical Physics Letters, **292**, 243 (1998).
- [3] For a review of Lie algebraic methods in molecular spectroscopy, see, F. Iachello, R.D. Levine, *Algebraic Theory of Molecules* (Oxford University Press, NY, 1995).
- [4] Y. Alhassid, F. Gürsey, F. Iachello, Ann. Phys. (NY), **148**, 346 (1983); Chem. Phys. Lett., **99**, 27 (1983); R.D. Levine, Chem. Phys. Lett., **95**, 87 (1983).
- [5] M. Ibrahim, <http://arXiv.org/abs/physics/0005036> .
- [6] O.S. van Roosmalen, *Algebraic Descriptions of Nuclear and Molecular Rotation-Vibration Spectra*, Ph.D. Thesis (University of Groningen, Netherlands, 1982).
- [7] C. Wulfman, Phys. Rev. A, **54**, R987 (1996).
- [8] R. Gilmore, C.M. Bowden, L.M. Narducci, Phys. Rev. A., **12**, 1019 (1975).
- [9] A.E.L. Dieperink, O. Scholten, F. Iachello, Phys. Rev. Lett., **44**, 1747 (1980).
- [10] The foundations of quantizing Lie groups to obtain algebraic quantum mechanics can be found in: B. Kostant, “Quantization and Unitary Representation,” Lecture Notes in Mathematics, Vol. 170 (Springer-Verlag, New York, 1970); J.-M. Soriau, “Structure des Systèmes Dynamiques,” (Dunod, Paris, 1970); alternatively one may use the more physical path integral approach in section 4 of [11].

- [11] For a review of the theory of coherent states, see, W. Zhang, D.H. Feng, R. Gilmore, *Rev. Mod. Phys.*, **62**, 867 (1990).
- [12] See, for example, F. Iachello, A. Arima, *The Interacting Boson Model* (Cambridge University Press, NY, 1987).
- [13] R. Gilmore, *Lie Groups, Lie Algebras, and Some of Their Applications* (John Wiley & Sons, NY, 1974).
- [14] See, for example, V.I. Arnold, *Mathematical Methods of Classical Mechanics* (Springer Verlag, NY, 1978), pp. 201ff.
- [15] B. Shao, N.R. Walet, R.D. Amado, *Phys. Rev. A*, **46**, 4037 (1992); B. Shao, N.R. Walet, R.D. Amado, *Phys. Rev. A*, **47**, 2064 (1993).
- [16] F. Iachello, S. Oss, *J. Chem. Phys.*, **104**, 6956 (1996).
- [17] E. Santopinto, R. Bijker, F. Iachello, *J. Math. Phys.*, **37**, 2674 (1996).
- [18] See, for example, S. Flügge, *Practical Quantum Mechanics*, Vol. I (Springer-Verlag, NY, 1971), pp. 107–110.
- [19] See, for example, M. Abramowitz, I. Stegun, *Handbook of Mathematical Functions* (Dover Publications, Inc., NY, 1965), pp. 255, 257, 508.
- [20] F. Iachello, S. Oss, *Phys. Rev. Lett.*, **666**, 2976 (1991).
- [21] F. Iachello, *Chem. Phys. Lett.*, **78**, 581 (1981); F. Iachello, R.D. Levine, *J. Chem. Phys.*, **77**, 3046 (1982).
- [22] See, for example, I.S. Gradshteyn, I.M. Ryzhik, *Table of Integrals, Series, and Products*, 5th ed. (Academic Press, NY, 1980), pp. 849, 850.

		Group Coordinates	Projective Coordinates
$\langle \hat{l} \rangle / N$	$\equiv l_{\text{cl}} \ (\tilde{l}_{\text{cl}})$	$q_x p_y - q_y p_x$	$2\Sigma(\tilde{q}_x \tilde{p}_y - \tilde{q}_y \tilde{p}_x)$
$\langle \hat{n} \rangle / N$	$\equiv n_{\text{cl}} \ (\tilde{n}_{\text{cl}})$	$\frac{1}{2}(q \cdot q + p \cdot p)$	$\Sigma(\tilde{q} \cdot \tilde{q} + \tilde{p} \cdot \tilde{p})$
$\langle \hat{n}^2 \rangle / N$		$n_{\text{cl}} + (N-1)(n_{\text{cl}})^2$	$\tilde{n}_{\text{cl}} + (N-1)(\tilde{n}_{\text{cl}})^2$
$\langle \hat{l}^2 \rangle / N$		$n_{\text{cl}} + (N-1)(l_{\text{cl}})^2$	$\tilde{n}_{\text{cl}} + (N-1)(\tilde{l}_{\text{cl}})^2$
$\langle \hat{W}^2 \rangle / N(N-1)$		$(l_{\text{cl}})^2 + 2(1 - n_{\text{cl}}) q \cdot q + \frac{2}{N-1}$	$(\tilde{l}_{\text{cl}})^2 + 4\Sigma^2 \tilde{q} \cdot \tilde{q} + \frac{2}{N-1}(\Sigma + \tilde{n}_{\text{cl}})$

Table 1: The coherent state limit of $U(3)$ operators in group and projective coordinates. In group coordinates we take the expectation in the state $\frac{1}{\sqrt{N!}}(\sqrt{1 - \alpha \cdot \alpha^*} \sigma^\dagger + \alpha \cdot \tau^\dagger)^N |0\rangle$ and let $\alpha = \frac{1}{\sqrt{2}}(q + ip)$ to reproduce the standard form of the angular momentum. In projective coordinates the unnormalized coherent state is parameterized by $\frac{1}{\sqrt{N!}}(\sigma^\dagger + \tilde{\alpha} \cdot \tau^\dagger)^N |0\rangle$. One must then divide the expectation by the normalization. We then let $\tilde{\alpha} = (\tilde{q} + i\tilde{p})$ to jibe with the standard definitions in the literature. Note that the ‘ q ’s and ‘ p ’s in each column are not the same. We have further made the abbreviation for projective coordinates: $\Sigma = \frac{1}{1 + (\tilde{q} \cdot \tilde{q} + \tilde{p} \cdot \tilde{p})}$.